

Precision predictions for Z' production at the LHC

Benjamin Fuks

*Physikalisches Institut, Albert-Ludwigs-Universität Freiburg,
Hermann-Herder-Straße 3, D-79106 Freiburg i.Br., Germany*



We present precision calculations of the transverse-momentum spectrum, the invariant-mass distribution and the total cross section for Z' production at hadron colliders. We implement joint resummation at the next-to-leading logarithmic accuracy and consistently match the obtained result with the pure perturbative result at the first order in the strong coupling constant. We confront our numerical predictions with the Monte Carlo generator PYTHIA and with a new implementation of extra neutral gauge bosons in the Monte Carlo generator MC@NLO. The impact of scale dependence is also studied.

1 Introduction

Despite its phenomenological success, the Standard Model (SM) of particle physics is believed to suffer from a variety of conceptual deficiencies. In particular, it provides no fundamental motivation why the strong and electroweak interactions should be described by three different gauge groups. Grand Unified Theories allow for a unification of these groups which is broken to the SM at higher scale. This yields one or several extra neutral gauge Z' -bosons exhibiting the existence of additional $U(1)$ symmetries¹. If the Z' -bosons couple to quarks and leptons not too weakly and if their mass is not too large, they will be produced at the Tevatron and the LHC and easily detected through their leptonic decay modes. The search for these particles occupies therefore an important place in the present experimental program.

When studying the transverse-momentum (p_T) distribution of a Z' -boson with an invariant mass M , it is convenient to separate the large- and small- p_T regions. For the large values of p_T , the use of the fixed-order perturbation theory is fully justified, since the perturbative series is controlled by a small expansion parameter, $\alpha_s(M^2)$, but in the small- p_T region, the coefficients of the perturbative expansion are enhanced by powers of large logarithmic terms, $\ln(M^2/p_T^2)$. As a consequence, the convergence of the perturbative series is spoiled as $p_T \rightarrow 0$. Furthermore,

when the initial partons have just enough energy to produce a Z' -boson, the mismatch between virtual corrections and phase-space suppressed real-gluon emission leads also to the appearance of large logarithmic terms $[\ln(1-z)/(1-z)]_+$, where $z = M^2/s$, s being the partonic centre-of-mass energy. However, the convolution of the partonic cross section with the steeply falling parton distributions enhances the threshold contributions even if the hadronic threshold is far from being reached and large corrections are expected.

Accurate calculations of p_T and invariant-mass distributions must then include soft-gluon resummation in order to obtain reliable perturbative predictions and properly take these logarithmic terms into account. We implement the joint resummation formalism² at the next-to-leading logarithmic (NLL) accuracy and consistently match the obtained result with the pure perturbative result at the first order in α_s ^a, which allows us to resum all the logarithms simultaneously. In addition, we compare our predictions⁴ with those of the next-to-leading order (NLO) Monte Carlo generator MC@NLO⁵ and of the leading-order (LO) Monte Carlo generator PYTHIA⁶, and study the impact of the scale dependence.

2 Theoretical framework

2.1 Considered Z' model

Ten-dimensional string theories with $E_8 \times E_8$ gauge symmetry are anomaly-free and contains chiral fermions as in the SM⁷. After compactification, this symmetry leads to an effective E_6 GUT group that can be broken further to⁸

$$E_6 \rightarrow SO(10) \times U(1)_\psi \rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi. \quad (1)$$

While the Z' -bosons related to the additional $U(1)_\psi$ and $U(1)_\chi$ symmetries can in general mix, we consider in this work only a TeV-scale Z_χ -boson and assume the Z_ψ to acquire its mass at much higher scales, as it is naturally the case in the hierarchy of symmetry breaking of Eq. (1).

2.2 Joint resummation formalism at the next-to-leading logarithmic level

In Mellin N -space, the resummed hadronic cross section for the hard scattering process

$$h_a h_b \rightarrow \gamma, Z, Z' \rightarrow l^+ l^- (M, p_T) + X, \quad (2)$$

where a lepton pair with invariant mass M and transverse momentum p_T is produced, factorizes

$$\frac{d^2\sigma^{(\text{res})}}{dM^2 dp_T^2}(N, b) = \sum_{a,b} f_{a/h_a}(N+1) f_{b/h_b}(N+1) \mathcal{H}_{ab}(N) \exp\{\mathcal{G}(N, b)\}. \quad (3)$$

$f_{a,b/h_{a,b}}$ are the N -moments of the universal distribution functions of partons a, b inside the hadrons $h_{a,b}$, the impact-parameter b is the variable conjugate to p_T through a Fourier transform and the dependence on the renormalization and factorization scales μ_R and μ_F has been removed for brevity. The process-dependent function $\mathcal{H}$ contains all the terms due to hard virtual corrections and collinear radiation while the process-independent Sudakov form factor $\mathcal{G}$ allows to resum the soft-collinear radiation, embodying the all-order dependence on the logarithms.

Once resummation has been achieved in N - and b -space, inverse transforms have to be performed in order to get back to the physical space. Since the resummed exponent contains singularities, the integration contours of the inverse transforms must avoid hitting any of these poles. The b -integration is performed by deforming the integration contour with a diversion into the complex b -space⁹, while the inverse Mellin transform is performed following a contour inspired by the Minimal Prescription¹⁰ and the Principal Value Resummation¹¹.

^aNote that next-to-next-to-leading order effects have recently been investigated as well³.




Figure 1: Transverse-momentum spectra (left), mass spectra (centre) and dependence of the total cross section on the common factorization/renormalization scale (right) in LO QCD (dot-dashed), NLO QCD (dashed) and after resummation (plain). The mass spectra have been normalized to the LO QCD prediction.

In order to keep the full information contained in the fixed-order calculation and to avoid possible double-counting of the logarithmically enhanced contributions, a matching procedure of the NLL resummed cross section to the fixed order result is performed through the formula

$$\frac{d^2\sigma}{dM^2 dp_T^2} = \frac{d^2\sigma^{(\text{F.O.})}}{dM^2 dp_T^2} + \oint_{C_N} \frac{dN}{2\pi i} \tau^{-N} \int \frac{b db}{2} J_0(b p_T) \left[\frac{d^2\sigma^{(\text{res})}}{dM^2 dp_T^2}(N, b) - \frac{d^2\sigma^{(\text{exp})}}{dM^2 dp_T^2}(N, b) \right]. \quad (4)$$

$J_0(x)$ is the 0th-order Bessel function, $d^2\sigma^{(\text{F.O.})}$ is the fixed-order perturbative result, $d^2\sigma^{(\text{res})}$ is the resummed cross section and $d^2\sigma^{(\text{exp})}$ is its truncation to $\mathcal{O}(\alpha_s)$.

3 Numerical results

In Fig. 1 (left), we show the p_T -spectrum (integrated over the invariant mass in the range from 900 to 1200 GeV) for Z' -bosons with a mass of 1 TeV produced at the LHC. As expected, the true NLO distribution (dashed) diverges as $p_T \rightarrow 0$ while resummation (plain) leads to a smooth turnover with a maximum at around 8 GeV. As can be seen from Fig. 1 (centre), the NLO K -factor (dashed) is increased further by the resummed contributions (plain), even if we are still relatively far from the production threshold. The scale uncertainty of the total cross section (integrated over all transverse momenta and over the invariant mass in the range from 900 to 1200 GeV) is shown in Fig. 1 (right). The LO QCD prediction (dot-dashed) does not give a reliable estimate of the theoretical error, since the NLO cross section (dashed) is considerably larger. At NLO, the factorization scale dependence is reduced as expected, but $\alpha_s(\mu_R)$ makes its appearance, so that an additional renormalization scale dependence is introduced. The total NLO scale dependence is reduced to 9% (vertical lines) once resummation is achieved (plain).

We confront in Fig. 2 the resummed results (plain) to the LO predictions obtained with PYTHIA's "power shower" (circles) and to the NLO predictions obtained with MC@NLO's parton shower (stars). The shaded band around the resummed predictions corresponds to the theoretical uncertainty induced through the simultaneous variation of the renormalization and factorization scales by a factor of two around the central scale M . The p_T -spectra (left) are, for all three calculations, no longer divergent. The PYTHIA prediction rises and falls rather steeply around its maximum at 3 GeV, whereas the MC@NLO and resummed predictions rise and fall more slowly around the peak at 8 GeV. The agreement between MC@NLO and joint resummation is impressive, in particular for a scale choice of $M/2$ (upper end of the shaded band). The correction factors for the mass spectra (right), which have been normalized to the LO QCD prediction, show only a very weak mass dependence. We have multiplied the PYTHIA




Figure 2: Mass (left) and transverse-momentum spectra (right) in PYTHIA (circles), in MC@NLO (stars), and after joint resummation (plain). The mass spectra have been normalized to the LO QCD prediction, and the renormalization and factorization scale uncertainties in the resummed predictions are indicated as shaded bands.

mass spectrum by hand with a global K -factor of 1.26. Otherwise, within the statistical error bars, the PYTHIA K -factor would just be unity, since the normalization of the total cross section is not changed by the parton shower. Again, the MC@NLO K -factor agrees almost perfectly with the one of resummation.

4 Conclusions

We have improved the theoretical predictions for the production of extra neutral gauge bosons at hadron colliders, which are currently based on the LO Monte Carlo generator PYTHIA, by implementing the Z' -bosons in the MC@NLO generator and by computing their differential and total cross sections in the framework of joint resummation. MC@NLO and resummation were found to be in excellent agreement for mass spectra and p_T spectra, while the PYTHIA predictions show significant shortcomings both in normalization and shape. The theoretical uncertainties from scale variations were found to be under good control. The implementation of our improved predictions in terms of the new MC@NLO generator or resummed K factors in the analysis chains of the Tevatron and LHC experiments should be straightforward and lead to more precise determinations or limits of the Z' -boson masses and/or couplings.

Acknowledgments

The author acknowledges the conference organizing committee for financial support.

References

1. P. Langacker, arXiv:0801.1345 [hep-ph], and references therein.
2. G. Bozzi, B. Fuks and M. Klasen, Nucl. Phys. B **794** (2008) 46.
3. C. Coriano, A. E. Faraggi and M. Guzzi, arXiv:0802.1792 [hep-ph].
4. B. Fuks, M. Klasen, F. Ledroit, Q. Li and J. Morel, Nucl. Phys. B **797** (2008) 322.
5. S. Frixione and B. R. Webber, JHEP **0206** (2002) 029.
6. T. Sjöstrand, S. Mrenna and P. Skands, JHEP **0605** (2006) 026.
7. M. B. Green and J. H. Schwarz, Phys. Lett. B **149** (1984) 117.
8. J. L. Hewett and T. G. Rizzo, Phys. Rept. **183** (1989) 193.
9. E. Laenen, G. Sterman and W. Vogelsang, Phys. Rev. Lett. **84** (2000) 4296.
10. S. Catani, M. L. Mangano, P. Nason and L. Trentadue, Nucl. Phys. B **478** (1996) 273.
11. H. Contopanagos and G. Sterman, Nucl. Phys. B **419** (1994) 77.